



**DR. BRIAN KLIKA & DR. ANDREW KIRKPATRICK
FLEXOR TENDON REPAIR ZONES 1-4 DELAYED MOBILIZATION
POST-OP THERAPY PROTOCOL**

The delayed mobilization program is appropriate for patients with significant soft tissue injury, crush injuries, young children under the age of 9, patients with cognitive disabilities or those who are otherwise unable to adhere to restrictions. In these instances, it is in the best interest of the patient to immobilize the hand in a cast for the first 3 weeks to avoid the risk of rupturing the repair.

Phase 1 – Maximum Protection (3-4½ weeks)

Goals for Phase 1

- Immobilize and protect repair
- Initiate ROM while protecting repair
- Minimize scar adhesions
- Edema control

Splint

- Dorsal blocking splint is fitted for continual wear in the following position:
 - Wrist: 20° extension, MP's: 60° flexion, IP's: full extension
 - Apply gutter splints as needed to maintain full extension
 - If there is a nerve repair, position PIP joint at 30° of flexion

ROM

- Initiate A/PROM to all digits within constraints of splint
- Begin with isolated passive flexion and extension of the MP, PIP, DIP followed by composite active and passive flexion/extension – 25 repetitions each
- If passive IP joint flexion is limited, a dynamic flexion component may be added to the DBS
- AROM to uninvolved joints

Scar Management

- Begin scar massage no sooner than 2 days after suture removal. Scar must be fully closed and have no scabbing present. Begin with light massage using lotion.
- Apply scar remodeling products as needed

Edema Control

- Digital level edema control can be initiated with Coban or compression sleeves
- Manual Edema Mobilization (MEM) with elevation

Modalities

- After 4 weeks, NMES may be added after 2-3 days of AROM.



**DR. BRIAN KLIKA & DR. ANDREW KIRKPATRICK
FLEXOR TENDON REPAIR ZONES 1-4 DELAYED MOBILIZATION
POST-OP THERAPY PROTOCOL**

Phase 2 – Maximize Active Range of Motion (4½ -6 weeks)

Goals for Phase 2	Precautions for Phase 2
• Continue to protect healing repair while maximizing active range of motion • Continue scar and edema control	• No composite passive wrist and digit extension

Splint

- Continue dorsal blocking splint between exercise sessions and at night

ROM

- Begin full AROM outside of the splint every 2 hours
- Continue phase 1 exercises
- Active tenodesis exercises: composite digit flexion with wrist extension followed by composite digit extension and wrist flexion
- Active composite wrist and finger exercises: wrist and finger flexion followed by wrist and finger extension
- Differential tendon glides: composite fist followed by MP extension and then IP extension

Manual Therapy

- Continue phase 1 scar and edema management



**DR. BRIAN KLIKA & DR. ANDREW KIRKPATRICK
FLEXOR TENDON REPAIR ZONES 1-4 DELAYED MOBILIZATION
POST-OP THERAPY PROTOCOL**

Phase 3 – Restore Strength and Functional Activity (6-12 weeks)

Goals for Phase 3	Precautions for Phase 3
• Restore full active and passive ROM • Regain strength • Return to ADL and full duty work	• Educate patient in no heavy lifting or use of the hand until 12-16 weeks post-op or physician consent

Splint

- Discontinue dorsal blocking splint
- If extrinsic flexor tightness is present, an extension splint for nighttime use may be appropriate

ROM

- PROM may be initiated to wrist and digits

Manual Therapy

- Continue phase 1 scar and edema management
- Ultrasound may be added for dense scar and/or limited tendon excursion

Strengthening

- At 8-10 weeks post-op, initiate wrist and hand strengthening

Work Conditioning

- After 10-12 weeks and with physician consent, a comprehensive work conditioning program for patients with high demand/heavy manual labor occupations may be appropriate