

DR. JOHN AWOWALE
ACL RECONSTRUCTION WITH MENISCUS REPAIR OR MICROFRACTURE
POST-OP THERAPY PROTOCOL

Phase 1 – Maximum Protection Phase (0-4 weeks)
(continued on next page)

Goals for Phase 1	Precautions for Phase 1	Criteria for Progression to Phase 2
• Protect graft and fixation • Minimize knee effusion and effects of immobilization • ROM 0-90°	• Avoid knee hyperextension during this phase greater than 10° • No kicking in pool for 12 weeks • No isolated resistance knee flexion for 12 weeks due to hamstring autograft • Avoid squatting and flexion on leg press beyond 90° until 4 months post-surgery	• Good PROM flexion/extension • Good quad set, SLR without extension lag • Minimal swelling/inflammation • Normal gait on level surfaces

Weight Bearing

- **0-2 Weeks:** Toe touch weight bearing (25%)
- **2-6 Weeks:** 50% weight bearing then advance to full weight bearing

Brace

- Brace 0-90° for ADLs until 6 weeks post-surgery

Range of Motion

- 0-90°
- Avoid hyperextension >10°

Manual Therapy

- Patellar mobility (superior, inferior, medial, lateral)
- Scar massage when incisions closed
- Gentle flexibility using deep tissue mobilization of surrounding tissues
- PROM/AROM knee flexion/extension, strong emphasis on full knee extension

Strengthening

- Quadriceps setting
 - Avoid knee hyperextension with quadriceps setting
- NMES to promote quad activation
- Multi-plane hip strengthening – start with SLR with brace locked in full extension until there is sufficient strength to prevent extension lag. Add resistance as tolerated
- 4-way hip strengthening, Standing TKE, mini step-ups, bridging, calf raises, mini squats
- Core strengthening
- For hamstring autograft:
 - Avoid isolated hamstring strengthening x 12 weeks
 - Heel slides to 90° only

Phase 1 – Maximum Protection Phase (0-4 weeks)

- For quadriceps and patellar tendon autograft:
 - Closed kinetic chain quadriceps strengthening activities as tolerated (wall sit, step ups, mini squats, leg press 90-30°)
 - Quadriceps isometrics at 60° and 90°
 - If available, aquatics for normalizing gait, weight bearing and strengthening
 - Stationary bike – initially for promotion of ROM – progress light resistance as tolerated
 - Hamstring curls

Aquatics

- Initiate aquatic therapy program when incisions are closed

Neuromuscular Control

- Proprioception on stable surface

Modalities

- Vaso pneumatic compression for edema management 2-3x/week (15-20 min)
- Cryotherapy at home, 3 x per day for 20 minutes each with knee elevated above heart
- NMES for quadriceps function
- Initiate blood flow restriction training after incisions are healed >3 weeks post-op – discuss with Dr. Awowale prior to initiation



DR. JOHN AWOWALE
ACL RECONSTRUCTION WITH MENISCUS REPAIR OR MICROFRACTURE
POST-OP THERAPY PROTOCOL

Phase 2 – Strengthening Phase (4-10 weeks)
(continued on next page)

Goals for Phase 2	Precautions for Phase 2	Criteria for Progression to Phase 3
• Restore normal gait with stair navigation after brace is discontinued at 6 weeks • Maintain full extension, progress toward full flexion ROM at 6+ weeks • Protect graft and fixation • Increase LE strength • Increase proprioception	• Avoid twisting and pivoting motions for 12 weeks • Avoidance of full body weight impact activity until able to pass functional testing	• No patellofemoral pain • Minimum of 120° knee flexion • Sufficient strength and proprioception to initiate running • Minimal swelling/inflammation

Brace

- Brace 0-90° for ADLs until 6 weeks post-surgery

Range of Motion

- Restore full ROM
- Maintain normal LE flexibility

Strengthening

- Stationary bike or elliptical – stair master as strength and gait allow
- Begin running in the pool (waist deep) or on an unweighted treadmill at 8 weeks.
 - Should have adequate strength, ROM, neuromuscular control, and limited swelling prior to initiation
- Bilateral gym strengthening with progression to unilateral as able (leg press, step-ups, hamstring curls, side-stepping, single leg squat, multi-directional lunges)
- Progress hip, hamstring and gastrocnemius strengthening
 - For hamstring autograft avoid isolated hamstring strengthening x 12 weeks
- Initiate knee flexion AROM using CKC strengthening with progression to OKC
- Core strengthening

Neuromuscular Control

- Proprioceptive drills progressing to on unstable surfaces
- Add dual tasking and reactive balance

Modalities

- Cryotherapy after activity
- Continue use of blood flow restriction training as need to build strength

Phase 2 – Strengthening Phase (4-10 weeks)

Testing to Advance to Phase 3 Protocol

- Functional strength testing to be scheduled before 10–12-week follow-up with MD. Appointment must be scheduled with Aurora BayCare Sports Physical Therapy at the 1110 Kepler location. Please contact physician office if unable to make this arrangement for alternative testing.
- Y-Balance testing within 6 cm of involved LE
- 3PQ isometric quadriceps testing (<25% difference)
- Single leg squat without display of knee valgus
- Recommend isokinetic test with anti-shear device at 12 weeks (14-16 weeks for hamstring tendon autograft procedures) to guide continued strengthening.

DR. JOHN AWOWALE
ACL RECONSTRUCTION WITH MENISCUS REPAIR OR MICROFRACTURE
POST-OP THERAPY PROTOCOL

Phase 3 – Strengthening and Plyometric Phase (10-16 weeks)

Goals for Phase 3	Precautions for Phase 3	Criteria for Progression to Phase 4
• Full ROM • Improve strength, endurance, and proprioception of the LE to prepare for sport activities • Normal running mechanics • Strength >70% of uninvolved LE with isokinetic evaluation	• Protect the patellofemoral Joint • Avoid overstressing the graft • Progressively increase resistance of hamstring (hamstring autograft)	• No significant swelling/inflammation. • Full, pain-free ROM • No evidence of patellofemoral joint irritation • Strength >70% of uninvolved lower extremity per isokinetic evaluation • Sufficient strength and proprioception to initiate agility activities • Normal running gait

Strengthening

- Stationary bike, elliptical, treadmill, may begin swimming
- Improve cardiovascular endurance
- Core strengthening
- Maintain LE flexibility – hamstring, quad, gastroc-soleus, ITB
- Initiate OKC knee extensions 90-30°, progress to eccentrics
- Unilateral gym strengthening program (single leg squats, eccentric leg press, lateral step-downs, advanced bridging, multi-directional lunges, CKC hamstring curls)
- If isokinetics available (with anti-shear device) – begin with mid-range speeds (120-240°/sec)
- Recommend isokinetic test with anti-shear device at 12 weeks (14-16 weeks for hamstring tendon autograft)
- **12+ weeks – for bone-patellar tendon-bone autograft:** Progress toward full weight bearing running - sagittal plane running, sub-maximal plyometrics
- **16+ weeks – hamstring tendon autograft:** Progress toward full weight bearing running - sagittal plane running, sub-maximal plyometrics

Neuromuscular Control

- Advanced proprioception on un-stable surfaces with perturbations and/or dual tasking, add sport specific balance tasks as able

Modalities

- Cryotherapy after activity

DR. JOHN AWOWALE
ACL RECONSTRUCTION WITH MENISCUS REPAIR OR MICROFRACTURE
POST-OP THERAPY PROTOCOL

Phase 4 – Advanced Strength and Advanced Plyometric Phase (4-6 months+)

Goals for Phase 4	Precautions for Phase 4	Criteria for Progression to Phase 5
- Symmetric performance of basic and sport specific agility drills - Single and 3 hop tests 85% of uninvolved LE - Quadriceps and hamstring strength at least 85% of uninvolved lower extremity per isokinetic strength test	None	- No patellofemoral or soft tissue pain or complaint - Necessary joint ROM, strength, endurance, and proprioception to safely return to work or athletics

Strengthening

- Continue advanced strengthening
- Agility progression including, but not limited to:
 - Side steps
 - Crossovers
 - Figure 8 running
 - Shuttle running
 - One leg and two leg jumping
 - Cutting
 - Acceleration/deceleration/sprints
 - Agility ladder drills
- Avoid** impact activities on unstable surfaces until >6 months post op or per conversation with Dr. Awowale with functional testing results.
- Promote adequate quad and hamstring strength
- Core and hip strengthening
- Begin building power in involved LE
- Assessment of running on treadmill
- Initiate sport-specific drills as appropriate

Neuromuscular Control

- Emphasize proper motor control
- Advanced proprioceptive drill examples:
 - Unsteady surface
 - Reactive balance
 - Deceleration control
 - Landing/take off drills
 - Perturbation training

Modalities

- As needed



DR. JOHN AWOWALE
ACL RECONSTRUCTION WITH MENISCUS REPAIR OR MICROFRACTURE
POST-OP THERAPY PROTOCOL

Phase 5 – Return to Activity Phase (6 months+)

Goals for Phase 5	Precautions for Phase 5
• Safe return to athletics/work • Maintenance of strength, endurance, proprioception • Patient education with regards to any possible limitations	• None

- Continue progression of activities from Phase 4
- Gradual return to sports participation
- Maintenance program for strength, endurance

Return to Function Testing: Aurora BayCare return to function for the lower extremity protocol to be used

- **6 months+: Return to function testing** per MD approval. Appointment must be scheduled with Aurora BayCare Sports Physical Therapy at the 1110 Kepler location. Please contact physician office if unable to make this arrangement for alternative testing.
- Criteria: Pain-free, full ROM, minimal joint effusion, isokinetic strength and functional testing at 90% compared to uninvolved, adequate knee control with sport and/or work specific tasks